




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Journal Pre-proof

Gargling with povidone iodine has a short-term inhibitory effect on SARS-Cov-2 in COVID-19 patients

Tetsuya Seikai, Ayato Takada, Akira Hasebe, Masahiro Kajihara, Kosuke Okuya, Tamaki Sekiguchi Yamada, Wataru Kakuguchi, Satoshi Konno, Yoichi Ohiro



PII: S0195-6701(22)00006-8

DOI: <https://doi.org/10.1016/j.jhin.2022.01.001>

Reference: YJHIN 6572

To appear in: *Journal of Hospital Infection*

Received Date: 28 December 2021

Revised Date: 7 January 2022

Accepted Date: 7 January 2022

Please cite this article as: Seikai T, Takada A, Hasebe A, Kajihara M, Okuya K, Sekiguchi Yamada T, Kakuguchi W, Konno S, Ohiro Y, Gargling with povidone iodine has a short-term inhibitory effect on SARS-Cov-2 in COVID-19 patients, *Journal of Hospital Infection*, <https://doi.org/10.1016/j.jhin.2022.01.001>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 The Healthcare Infection Society. Published by Elsevier Ltd. All rights reserved.

**Gargling with povidone iodine has a short-term inhibitory effect on SARS-Cov-2 in COVID-19 patients**

**Running title: Gargling effect with PVP-I on SARS-Cov-2**

Tetsuya Seikai<sup>a</sup>, Ayato Takada<sup>b,c</sup>, Akira Hasebe<sup>d</sup>, Masahiro Kajihara<sup>b</sup>, Kosuke Okuya<sup>b</sup>, Tamaki Sekiguchi (Yamada)<sup>a</sup>, Wataru Kakuguchi<sup>a</sup>, Satoshi Konno<sup>e</sup> and Yoichi Ohiro<sup>a\*</sup>

<sup>a</sup> Department of Oral and Maxillofacial Surgery, Faculty of Dental Medicine and Graduate School of Dental Medicine, Hokkaido University, Kita-13, Nishi-7, Kita-ku, Sapporo, 060-8586, Japan

<sup>b</sup> Division of Global Epidemiology, International Institute for Zoonosis Control, Hokkaido University, Kita-20, Nishi-10, Kita-ku, Sapporo, 001-0020, Japan

<sup>c</sup> International Collaboration Unit, International Institute for Zoonosis Control, Hokkaido University, Kita-20, Nishi-10, Kita-ku, Sapporo, 001-0020, Japan

<sup>d</sup> Department of Oral Molecular Microbiology, Faculty of Dental Medicine and Graduate School of Dental Medicine, Hokkaido University, Kita-13, Nishi-7, Kita-ku, Sapporo, 060-8586, Japan

<sup>e</sup> Department of Respiratory Medicine, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Kita-15, Nishi-7, Kita-ku, Sapporo, 060-8638, Japan

\*Corresponding author. Mailing address:

Yoichi Ohiro DDS Ph.D.

Department of Oral and Maxillofacial Surgery, Faculty of Dental Medicine and Graduate School of Dental Medicine, Hokkaido University  
Kita-13, Nishi-7, Kita-ku, Sapporo, 060-8586, Japan

Phone: +81-11-706-4283

Fax: +81-11-706-4283

E-mail: [yohiro@den.hokudai.ac.jp](mailto:yohiro@den.hokudai.ac.jp)

1 It is known that povidone iodine (PVP-I) solutions have virucidal action against severe  
2 acute respiratory syndrome coronavirus (SARS-CoV) *in vitro* [1,2,3]. In this study, the  
3 saliva of coronavirus disease 2019 (COVID-19) patients was collected up to 2 hours after  
4 PVP-I gargling and the dynamics of SARS-CoV-2 infectivity in saliva were assessed by  
5 a real-time reverse transcription–polymerase chain reaction (rRT–PCR) and  
6 determination of the infectious viral load.

7 Patients (aged  $\geq 20$  years) who had symptoms indicative of SARS-CoV-2 infection  
8 within the last 7 days or asymptomatic patients with a cycle threshold  $< 40$  for SARS-  
9 CoV-2 ribonucleic acid (RNA), as determined by rRT–PCR of saliva, were included in  
10 this study (n=35). Patients who had an iodine allergy or thyroid disease were excluded.  
11 This study was approved by the Institutional Review Board (Hokkaido University  
12 Hospital Division of Clinical Research Administration Number: 020-0111), and written  
13 informed consent was obtained from all participants.

14 Baseline saliva samples were collected prior to intervention with iodine. Then, patients  
15 rinsed their mouths for 20 seconds with 20 mL of PVP-I gargle solution (Meiji Co., Ltd,  
16 Tokyo, Japan), which was diluted 15 times with water. Patients repeated gargling with  
17 PVP-I three times, then rinsed their mouths with water. After gargling, saliva was  
18 collected at four time points: immediately after gargling, and 30, 60, and 120 minutes  
19 (min) later. Patients collected saliva samples themselves by spitting into a sterile cup (PP  
20 Screw cup 50; ASIAKIZAI Co., Tokyo, Japan). Viral RNA was quantified in the samples  
21 by RT–PCR and the virus was titrated in cultured cells.

22 For RT–PCR, 200  $\mu$ L of saliva was added to 600  $\mu$ L of PBS, mixed vigorously, then  
23 centrifuged at  $20,000 \times g$  for 5 min at 4°C, and 140  $\mu$ L of the supernatant was used as the  
24 sample. rRT-PCR was conducted in accordance with the manual for the Detection of

Pathogen 2019-nCoV Ver. 2.9.1. (<https://www.niid.go.jp/niid/images/lab-manual/2019-nCoV20200319.pdf>). Total RNA was extracted using the QIAamp Viral RNA Mini Kit (QIAGEN, Hilden, Germany) and rRT-PCR was performed using the QuantiTect Probe RT-PCR Kit (QIAGEN) in the QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA). The sequences of the primers and TaqMan probe used for detection of the SARS-CoV-2 genome were as follows: forward primer (NIID\_2019 - nCoV\_N\_F2, 5' AAATTTTGGGGACCAGGAAC 3'), reverse primer (NIID\_2019 - nCoV\_N\_R2, 5' TGGCAGCTGTGTAGGTCAAC 3'), TaqMan probe (NIID\_2019 - nCoV\_N\_P2, 5' FAM-ATGTCGCGCATTGGCATGGA-BHQ 3').

Viral titers were determined as the 50% tissue culture infective dose (TCID<sub>50</sub>) of the virus. Vero E6 cells expressing the type II transmembrane serine protease (Vero-TMPRSS2) [4] were seeded into 96-well plates and incubated with a serial dilution of patient saliva. Three days later, cytopathic effects were examined. The samples in which infectious SARS-CoV-2 was detected before PVP-I gargling (i.e.,  $> 10 \times \text{TCID}_{50}$  of the virus) were targeted in this study.

Of a total of 35 COVID-19-positive patients, 24 were excluded from the study because they had undetectable SARS-CoV-2 RNA or a viral titer of  $< 10 \times \text{TCID}_{50}$  in their baseline saliva sample. Thus, 11 patients were analyzed in this study. The average viral RNA copies and viral titers were compared at each time point using the Wilcoxon rank sum test. The p-value threshold for significance was set at  $< 0.05$ .

Figure 1A shows the change in viral RNA copies (log<sub>10</sub> copies/mL) after PVP-I gargling. A statistically significant decrease in viral RNA was observed in the samples taken immediately after gargling and at 30 and 60 min, compared with before gargling.

Figure 1B shows the change in viral titer ( $\log_{10}$  TCID<sub>50</sub>/mL) after PVP-I gargling. A statistically significant decrease was observed in viral titer immediately after gargling and at 60 min, compared with before gargling. The viral titers in the samples at 30 min showed no significant difference ( $p = 0.055$ ), but the median value was lower compared with the samples taken before gargling.

In conclusion, viral copies and titers were significantly decreased 60 min after gargling. The reason for the temporary increase in viral titer at 30 min may become clear as the number of examined cases increases. However, importantly, our data indicated that PVP-I gargling effectively suppressed SARS-CoV-2 infectivity in saliva for 60 min. The application of PVP-I may be an effective measure to reduce the infection risk in situations such as during dental treatment and oral examination by physicians.

We recognize a limitation of our study is that this study was performed under simple conditions to minimize the risk of infection, and was carried out without a control group for gargling just with water. Despite the limitation, our findings support the use of PVP-I gargling in preventing infections via saliva over a short period.

## Acknowledgements

This work was supported by the accommodation medical treatment team of Sapporo Public Health Office, we are especially grateful to Dr. Akino and R.N. Mizuta.

## Conflicts of Interest

None declared.

## Funding



1 None.

Journal Pre-proof

## References

- [1] Anderson DE, Sivalingam V, Kang AEZ, Ananthanarayanan A, Arumugam H, Jenkins TM. Povidone-iodine demonstrates rapid in vitro virucidal activity against SARS-CoV-2, the virus causing COVID-19 disease. *Infect Dis Ther* 2020;9:669–75.
- [2] Bidra AS, Pelletier JS, Westover JB, Frank S, Brown SM, Tessema B. Rapid In-Vitro Inactivation of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Using Povidone-Iodine Oral Antiseptic Rinse. *Prosthodont* 2020;29:529–33.
- [3] Shet M, Hong R, Igo D, Cataldo M, Bhaskar S. In Vitro Evaluation of the Virucidal Activity of Different Povidone-Iodine Formulations Against Murine and Human Coronaviruses. *Infect Dis Ther* 2021;28:1-14.
- [4] Sasaki M, Uemura K, Sato S, Toba S, Sanaki T, Maenaka K, et al. SARS-CoV-2 variants with mutations at the S1/S2 cleavage site are generated in vitro during propagation in TMPRSS2-deficient cells. *PLoS Pathog* 2021;17:1–17.





**Figure Legend**

Figure 1. Changes in the SARS-COV-2 RNA level (A) and viral titer (B) in saliva samples before and after povidone iodine gargling

Box plots (median, interquartile range, 5th and 95th percentile). SARS-CoV-2: severe acute respiratory syndrome coronavirus 2. TCID<sub>50</sub>: tissue culture infectious dose. RNA: ribonucleic acid. Before: before gargling, Immediately after, immediately after gargling, min: minutes after gargling.